

Original Article

Systemic Hypertension among Adults in a Peri-Urban Community in Delta State Nigeria

Ochei O^{1,4*}, Aigbe F I^{2,3}, Utake O E⁴, Olamide O⁵, Oluonye H C⁵, Onike C⁵, Omosivwe E O⁵

¹ Department of Community Medicine, Delta State University, Abraka Nigeria. ² Department of Internal Medicine, Delta State University, Abraka Nigeria. ³ Department of Internal Medicine, Delta State University Teaching Hospital, Oghara Nigeria. ⁴ Department of Community Medicine, Delta State University Teaching Hospital, Oghara Nigeria. ⁵ Delta State University Teaching Hospital, Oghara, Nigeria.

Article History

Submitted: 15 September 2022; Accepted: 10 November 2022; Published: 20 December 2022

*Correspondence: Oboratare Ochei. Email: droboratare@gmail.com

ABSTRACT

The upsurge in the prevalence of systemic hypertension and its complications in urban and rural communities in Nigeria cannot be ignored. Periodic surveys on the trend of hypertension in communities are important in the prevention and treatment of the disease. This study was conducted to determine the prevalence, prevailing risk factors, and the level of knowledge of the disease, treatment types, and adherence to antihypertensive medications in a community in Nigeria. A cross-sectional design and a multistage sampling method were used in this study. The JNC7 guideline was used to decide the cut-off value of hypertension. From the data obtained, 38% were hypertensive of which 27.8% were previously diagnosed. The level of knowledge of hypertension and controlled blood pressure was low in this study. There was a statistically significant difference between blood pressure and the following risk factors: age, sex, marital status, exercise, BMI, and waist circumference. While the risk of developing hypertension was 2.21 times higher in cigarette smokers than in non-smokers. Participants who engaged in physical exercise had a 47% lower risk of developing hypertension when compared to those who were inactive; the odds ratio = 0.53. Lifestyle modifications and health promotion can prevent hypertension.

Keywords: Adults, Systemic Hypertension, Urban Nigeria.

INTRODUCTION

Systemic hypertension or High Blood Pressure is a significant risk factor for the development of Cardiovascular Diseases¹. In many countries of the world, hypertension is defined by a systolic and diastolic blood pressure of ≥ 140 mmHg and ≥ 90 mmHg respectively, but in the current guideline of the American College of Cardiology/American Heart Association (ACC/AHA), hypertension is

defined as systolic blood pressure/diastolic blood pressure of $\geq 130/80$ mmHg².

Its incidence is on the rise and currently affects a large percentage of adults worldwide. Approximately, a billion adults had hypertension in 2015 and 13.5% of annual deaths are ascribed to hypertension and its complications^{1,2,3}. Every year, complications from hypertension account for 9.4

Article Access



Website: www.wjmb.com

 10.5281/zenodo.7749310

How to cite this article

Ochei O, Aigbe F I, Utake O E, Olamide O, Oluonye H C, Onike C, Omosivwe E O. Systemic Hypertension Among Adults in a Peri-Urban Community in Delta State Nigeria. *West J Med & Biomed Sci.* 2022;3(3-4):103-113. DOI:10.5281/zenodo.7749310

million deaths, 45% of deaths due to heart diseases, and 51% of deaths due to stroke^{4,5,6}. The prevalence rate in low-income and high-income countries is approximately 31.5% and 28.5%⁷. In recent studies in Africa, about 20–46% of adults were reported to be hypertensive^{4,8-10}. Its prevalence varies with the study population, type of measuring instruments, and cut-off values used for defining hypertension. In Nigeria, prevalence studies in rural populations range from 13.5%–46.4%, urban, 8.1%–42.0%^{8,11-13}. Adeloje *et. al.*, 2015; obtained a prevalence of 30.6% among urban and 26.4% among rural Nigerians from a systematic review of population-based studies from 1990 to 2013¹⁴.

The global increase in the prevalence of hypertension has been linked to various factors, including aging, obesity, race, unhealthy diet, harmful use of alcohol, cigarette smoking, lack of physical activity, the stress of urbanization, and underlying diseases¹⁴⁻¹⁷. The higher the pressure in blood vessels, the harder the heart has to work to perfuse the tissues. Persistently uncontrolled blood pressure can lead to various manifestations of hypertensive heart disease and eventually culminate in heart failure. The increased blood pressure may also result in arteriosclerosis ultimately leading to arteriosclerotic vascular disease manifesting clinically as coronary artery disease, peripheral artery disease, and cerebrovascular disease. Hypertension can also lead to renal impairment, visual disturbance, and cognitive impairment¹⁴. It has a negative impact on the quality of life of affected individuals and on countries' health expenditures, consequently, negatively impacting the economy. In some low- and middle-income countries, health expenditure on cardiovascular diseases alone account for 20% of total health expenditure². Premature deaths, disability, psychosocial disorders, loss of income, and healthcare costs due to hypertension take a toll on families, communities, and nations' finances. In poor countries, many people are unable to seek standard care for hypertension because of its cost. Households often spend a substantial proportion of their income on hospitalization and care following complications of

hypertension².

Epidemiological studies show an inverse relationship between regular blood pressure checks and organ damage¹⁸. Since pre-hypertensives patients have been documented to have an increased risk of organ damage (compared to normotensive patients) it is therefore apparent that early detection, regular blood pressure check and adherence to treatment are likely to decrease the morbidity and mortality associated with hypertension¹⁹. Unfortunately, studies show that only 50% of those who are hypertensive are aware of their condition and less than 50% of those who are aware are on treatment, and less than 50% of those on treatment are receiving adequate management, a situation that has been referred to as the rule of halves^{20,21}. In Nigeria, 5.2% to 29.4% of hypertensive are aware of their condition^{15,23-24}. The low visibility of this condition and World Health Organization (WHO) advocacy for increased surveillance of non-communicable diseases have prompted this community survey on hypertension. Therefore, studies that create awareness, and measure community blood pressure, should be continually encouraged.

In this study, we determined the prevalence of hypertension, detected hypertensives who are not aware of their condition, assessed the level of knowledge of hypertension, risk factors, treatment types, and compliance to Antihypertensives among adults in an urban community in Nigeria. The study aimed at increasing the level of community awareness through screening. Information from this study will serve as important data for comparison with present/future data to track changes. It will also help in planning interventional and local research for the prevention and control of hypertension.

MATERIALS AND METHODS

The study was conducted in Oghara, a community in Delta State, located in the South-South geopolitical zone of Nigeria. It lies between longitude 6.1002°E and latitude 5.5867°N. The settlement is made up of five wards. It has a tertiary health facility, a polytechnic, a naval institution, gas stations, and

numerous gas storage tanks. The majority of the population are farmers, petty traders, and artisans. However, a sizeable proportion is engaged by organized public and private sectors.

A cross-sectional study design was used and the study population was adults aged 18 years and above who had resided in the community for a minimum of one year. Pregnant women were excluded from the study. The sample size was 255, estimated using the formula for cross-sectional studies. A previous study in the community reported a prevalence of 21% [25]. A multistage sampling method was used; in the first stage, three of the five political wards were selected by balloting. In the second stage, the calculated sample size of 255 was divided equally among the three selected wards, hence, 85 adults were recruited from each selected ward. The residential houses in the selected wards were numbered and divided by the sample size of 85 to obtain the sampling interval. The “computed sampling interval” was used to select a residential house from the study area; one adult above the age of 18 years was picked from each of the selected houses and informed consent was obtained.

The quantitative method was used to obtain data. The study tools include the following: a structured questionnaire, weighing scale, mercury sphygmomanometer, stethoscope, and calibrated rule. The questionnaire was interviewer-administered with the following variables: sociodemographic characteristics, knowledge of hypertension and risk factors, history of hypertension, whether or not presently on antihypertensive drugs, routine blood pressure check, and adherence to antihypertensives. Participants were also asked if they indulged in physical exercise, alcohol, and smoking cigarette. The number of fruits and vegetables consumed per day was also determined.

Participants who took at least 1 cigarette per day in the last year prior to the survey were classified as smokers while those who had never smoked or quit smoking for at least one year were classified as non-smokers¹⁸. Participants were also classified into groups based on the units of alcohol consumed in a week: occasional drinkers (< 1 unit), mild drinkers (1-<14units), moderate drinkers (14-21units), and heavy drinkers

(>21units)²⁵. One standard unit of alcohol is half a bottle of beer or a glass of alcoholic wine or one shot of gin/whisky. Physical exercise was classified based on work, leisure, and sports. Participants whose daily physical activities are either sedentary or light are categorized as physically inactive while those whose physical activities can be described as moderate and vigorous are classified as physically active²⁵. “Adequate fruit and vegetable consumption was defined as consuming 5 or more servings of fruits and/or vegetables per day²⁶.”

The weight, height, waist circumference, and blood pressure of participants were taken and the body mass index (BMI) was calculated, as weight (kg) divided by the square of height (m²). The weight was taken using a standard weighing scale. Height was taken using a calibrated rule. Blood pressure was measured with participants comfortably sitting, back well-supported, elbow resting on the table and legs uncrossed resting on the floor.

The diagnosis of hypertension was made when a participant is on treatment for hypertension or detection of systolic and/or diastolic blood pressure ($\geq 140/90$ mmHg) during the survey on two occasions using a standardized mercury sphygmomanometer. The seventh Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC7) 2003 guideline was used to decide the cut-off value²⁷. Participants were told their readings and counselled.

Data collected were analysed using the computer software SPSS version 26. Age, BMI, waist circumferences, and systolic and diastolic blood pressure measurements were summarised using mean, median, standard deviation, and range. Most variables were summarised using frequencies and percentages. The explanatory variables were sex, age, level of education, occupation, marital status, fruits and vegetable consumption; cigarette use, alcohol use, BMI, and waist circumference. The outcome variables were participants' hypertension status and knowledge of hypertension. Knowledge was assessed with twenty-two questions and a correct answer was scored one mark. Persons who scored 70% and above

were classified as having good knowledge; persons with scores of 50% to 69.9% had fair knowledge and those with scores less than 50% had poor knowledge. The differences between means in hypertensive and normotensive were compared using the independent t-test while the chi-square test was used to test for associations between hypertension status and risk factors. Odds ratios and confidence intervals were calculated if the explanatory and outcome variables measured were qualitative.

RESULTS

A total of 255 adults with 123 (48.2%) males and 132 (51.8%) females participated in the study. Their mean age was 41.73 ± 18.3 years, and the majority of the participants had secondary education 110 (43.1%), followed by tertiary 98 (38.4%). There were more married persons 118 (46.3%), than singles 107 (42.0%). About 10 (3.8%) of the participants were underweight, 128 (50.2%) had normal weight, 93 (36.5%) were overweight (table 1).

Table 2, shows the knowledge of the participants regarding hypertension: most had poor knowledge of hypertension 121(47.5%), followed by fair knowledge 71 (27.8%), while 63 (24.7%) had good knowledge. About 116 (45.5%) of the participants knew that cigarette smoking is a risk factor for hypertension, 134 (52.5%) knew that obesity is a risk factor, and 144 (56.5%) knew age as a risk factor, and race 63 (24.7%). Only about 159 (62.4%) and 68 (26.7%) knew that stroke and renal failure are complications of hypertension respectively.

The proportion of participants who had hypertension was 97 (38%) of which 71 (27.8%) were previously diagnosed, while 26 (10.2%) were diagnosed during the survey. Of the 97 hypertensives, 53 (54.6%) checked their blood pressure periodically. There were 118 (46.3%) normotensives and 40 (15.7%) pre-hypertensives, of which 26 (16.5%) checked their blood pressures periodically. The proportion of previously diagnosed hypertensives on antihypertensive medications was 57 (57/71, 80.3%). Fourteen of the hypertensives

(14/71=19.7%) were on herbal drugs only and 4 (4/71, 5.6%) were on both antihypertensive and herbal drugs. Only 39 (39/71, 54.9%) of the previously diagnosed hypertensives were compliant with their drugs and only 11(11/71, 15.5%) had their blood pressures under control at the time of the survey (table 3). There was statistically significant difference between age, sex, marital status and hypertension: more in males (46.3%) than females (30.3%), p -value = 0.008; more in elderly (78.3%) compared to middle-aged (56.9%) and young (18.5%), p -value = 0.0001; more in married (55.9%) than singles (15.9%), p -value = 0.0001. There was also a statistically significant association between exercise, BMI, waist circumference, and hypertension. Hypertensive mean waist circumference was 89 ± 10 cm while normotensive mean waist circumference was 82 ± 13 cm, p -value = 0.001. Hypertensive mean BMI was 25.8 ± 4.0 kgm² while normotensive mean BMI was 24 ± 4.0 kgm², p -value = 0.003 (table 4 and 5). Those who exercised (30.2%) had less hypertension than those who did not (44.8%), p -value = 0.017.

There was no statistically significant difference between alcohol, cigarette smoking, fruit and vegetable consumption, and hypertension. However, cigarette smokers showed a risk of 2.21 times more than non-smokers. Participants who engaged in physical exercises had a 47% lower risk of developing hypertension when compared to those that were inactive; the odds ratio = 0.533. Males showed a risk of 1.98 times of developing hypertension than females (table 4 and 5).

Table 1: Characteristics of adults in the study population (Oghara)

Variables	Frequency n=255 (%)
Age	
18-45	151 (59.3)
46-59	58 (22.7)
60-101	46 (18.0)
Gender	
Male	123 (48.2)
Female	132 (51.8)
Level of education	
None	12 (4.7)
Primary	35 (13.7)
Secondary	110 (43.1)
Tertiary	98 (38.4)
Marital status	
Married	118 (46.3)
Single	107 (42.0)
Widowed	14 (5.5)
Separated	8 (3.1)
Cohabiting	5 (2.0)
Divorced	3 (1.2)
Occupation	
Unemployed	67 (26.3)
Self-employed	122 (47.8)
Civil Servant	41 (16.1)
Others	25 (9.8)
BMI (Kg/m²)	
Underweight	10 (3.8)
Normal weight	128 (50.2)
Overweight	93 (36.5)
Obese	18 (7.1)
Grossly obese	6 (2.4)
Waist circumference	
Male	
<120cm	123 (48.2)
Female	
<88cm	89 (34.9)
≥88cm	43 (16.9)

Median BMI = 24.6(Kg/m²), Range 13.8-36.6(Kg/m²); Median Waist circumference 83cm, Range 60-112 cm.

Table 3: Blood pressure status of participants

Variables	Frequency, N=255 (%)
Blood Pressure Status	
Normotensive	118 (46.3)
Hypertensive (<i>unaware</i>)	26 (10.2)
Hypertensive (<i>aware</i>)	71 (27.8)
Prehypertensive (<i>unaware</i>)	40 (15.7)
Hypertensives that are aware (n=71)	
BP under control	11 (15.5)
On antihypertensive drugs	57 (80.3)
On herbal drugs only	14 (19.7)
On both antihypertensive and herbal drugs	4 (5.6)
Compliant with antihypertensive drugs	39 (54.9)
Blood pressure readings at the survey (mmHg)²⁷	
Normal (≤120mmHg systolic & ≤80mmHg diastolic)	129 (50.6)
High Normal (≤120-139 systolic and >80-90 diastolic)	62 (24.3)
Grade 1 (≥140-159 systolic and >90-100 diastolic)	51 (20.0)
Grade 2 (≥160mmHg systolic and >100mmHg diastolic)	13 (5.1)
*Duration of Hypertension (n =71)	
<5years	34 (47.8)
5-10 years	25 (35.2)
≥ 10 years	2 (2.8)
Practice Periodic Blood Pressure Check	
Hypertensive (<i>aware</i>)	53 (53/71 =74.6)
Normotensives + Prehypertensive (<i>unaware</i>)	26 (26/158 = 54.9)

Median Systolic Blood Pressure = 129mmHg, Range=90-178 mmHg; Median Diastolic Blood Pressure = 80mmHg, Range = 60-120mmHg

Table 2: Knowledge of hypertension of participants

Variables	Frequency, n=255 (%)
General knowledge of hypertension	
Good	63 (24.7)
Fair	71 (27.8)
Poor	121(47.5)
Know the definition of hypertension	168 (65.9)
Know the risk factors of hypertension	
Race	63 (24.7)
Sedentary lifestyle	89 (34.9)
Family history	99 (38.8)
High fat diet	92 (36.1)
Increased salt intake	116 (45.5)
Alcohol intake	120 (47.1)
Cigarette smoking	116 (45.5)
Obesity	134 (52.5)
Age	144 (56.5)
Think hypertension can be cured	
Yes	53 (20.8)
No	149 (58.4)
I don't know	53 (20.8)
Agreed that hypertension can be prevented	
Yes	167 (65.5)
No	38 (14.9)
I don't know	49 (19.2)
Agreed that hypertension has orthodox treatment	
Yes	
No	191 (74.9)
I don't know	31 (12.2)
Knows complications that can arise	32 (12.5)
Stroke	159 (62.4)
Heart attack	141(55.3)
Renal failure	68 (26.7)
Blindness	30 (11.8)
Leg ulcer	23 (9.0)
Diabetes	50 (19.6)

Table 4: Relationship between sociodemographic characteristics and hypertensive status of participants

Variable	Present	Hypertension Absent	Total
Age (years)			
Young (18-45)	28 (18.5%)	123 (81.5%)	151 (100.0%)
Middle (46-59)	33 (56.9%)	25 (43.1%)	58 (100.0%)
Elderly (60-101)	36 (78.3%)	10 (21.7%)	46 (100.0%)
$\chi^2 = 64.676$, p-value = 0.0001			
Sex			
Male	57 (46.3%)	66 (53.7%)	123 (100.0%)
Female	40 (30.3%)	92 (69.7%)	132 (100.0%)
$\chi^2 = 6.949$, p-value = 0.008			
OR= 1.98(CI=1.19-3.32)			
Marital Status			
Cohabiting	1 (20%)	4 (80%)	5 (100.0%)
Divorced	2 (66.7%)	1 (33.3%)	3 (100.0%)
Married	66 (55.9%)	52 (44.1%)	118 (100.0%)
Single	17 (15.9%)	90 (84.1%)	107 (100.0%)
Separated	4 (50.0%)	4 (50.0%)	8 (100.0%)
Widowed	7 (50.0%)	7 (50.0%)	14 (100.0%)
$\chi^2 = 41.373$, p-value = 0.0001			
Level of Education			
None	4 (33.3%)	8 (66.7%)	12 (100.0%)
Primary	19 (54.3%)	16 (45.7%)	35 (100.0%)
Secondary	42 (38.2%)	68 (61.8%)	110 (100.0%)
Tertiary	32 (32.7%)	66 (67.3%)	98 (100.0%)
$\chi^2 = 5.240$, p-value = 0.155			
Occupation			
Civil Servant	18 (43.9%)	23 (56.1%)	41 (100.0%)
Others	9 (36.0%)	16 (64.0%)	25 (100.0%)
Self-employed	49 (40.2%)	73 (59.8%)	122 (100.0%)
Unemployed	21 (31.3%)	46 (68.7%)	67 (100.0%)
$\chi^2 = 2.150$, p-value = 0.542			
Family History of Hypertension			
Yes	41 (44.1)	52 (55.9)	93 (100.0%)
No	32 (29.6)	76 (70.4)	108 (100.0%)
I Don't Know	24 (44.4)	30 (55.6)	54 (100.0%)
$\chi^2 = 5.623$, p-value = 0.060			
OR= Odds Ratio, CI= Confidence Interval			

Table 5: Relationship between alcohol intake, cigarette smoking, intake of fruits/vegetables, and excess fried food, balanced diet, and hypertensive status of participants

Variable	Hypertension		Total
	Present (n=97)	Absent (n=158)	
Alcohol Intake			
None	67 (39.6%)	102 (60.4%)	169 (100.0%)
Occasional users	16 (31.4%)	35 (68.6%)	51 (100.0%)
Mild users	8 (33.3%)	16 (66.7%)	24 (100.0%)
Moderate users	6 (54.5%)	5 (45.5%)	11 (100.0%)
$X^2 = 2.64$, p-value = 0.450			
Cigarette Smoking			
At least 1 cigarette/day	9 (56.3%)	7 (43.7%)	16 (100%)
Never smoke or quit	88 (36.8%)	151 (63.2%)	239 (100.0.0%)
$X^2 = 2.40$, p-value = 0.121			
OR= 2.21(CI=0.79-6.13)			
Physical Exercise			
Yes	36 (30.2%)	83 (69.8%)	119 (100.0%)
No	61 (44.8%)	75 (55.2%)	136 (100.0.0%)
$X^2 = 5.74$, p-value = 0.017			
OR= 0.533(CI=0.318-0.894)			
Excess Intake of Fried/Junk Food			
Yes			
No	73 (39.0%)	114 (61.0%)	187 (100.0%)
	24 (35.3%)	44 (64.7%)	68 (100.0%)
$X^2 = 7.65$, p-value = 0.382			
OR= 1.17 (CI=0.478-1.518)			
Intake of Fruits and Vegetables			
Appropriate	49 (43.0%)	65 (57.0%)	114 (100.0%)
Inappropriate	48 (42.1%)	93 (57.9%)	141 (100.0%)
$X^2 = 2.14$, p-value = 0.144			
BMI Status			
Underweight	3 (30.0%)	7 (70.0%)	10 (100.0%)
Normal weight	39 (30.5%)	89 (69.5%)	128 (100.0%)
Overweight	42 (45.2%)	51 (54.8%)	93 (100.0%)
Obese	11 (61.1%)	7 (38.9%)	18 (100.0%)
Grossly obese	2 (33.3%)	4 (66.7%)	6 (100.0%)
$X^2 = 9.51$, p-value = 0.050			
BMI (Mean±2SD)	25.8±4.0kg/m ²	24.2±4.0kg/m ²	
t = 3.02, p-value=0.003			
WC (women)			
< 88cm	22 (24.7%)	67 (75.3%)	89 (100.0%)
≥ 88-120cm	18 (41.9%)	25 (88.1%)	43 (100.0%)
$X^2 = 4.03$, p-value = 0.045			
WC women(Mean±2SD)	89.1±13.6cm	82±10.2cm	
t = 3.255, p-value=0.001			

BMI= Body Mass Index, WC = Waist Circumference.

DISCUSSION

We recorded an 11% increase in the prevalence of hypertension compared with the previous survey in the community, using the same case definition^{25,27}. The prevalence in this study was also higher than some previous studies conducted in Nigeria and internationally^{13,28-37}. It is, however, similar to values obtained by Daniel et al 2013²⁴; among urban slum dwellers in Lagos; Osunkwo et al 2020³⁸; among adults in Benue State, and within the range obtained from a nationwide study and a systematic review^{37,39}. The reasons for the higher prevalence rate compared to previous ones may be due to the progressive changes in the trend of the disease.

The proportion with pre-hypertension was observed to be three of every twenty persons. In the nationwide study conducted by Odili et al 2020³⁹; one fifth had pre-hypertension, while at Ibadan about one-third were pre-hypertensive¹³. The finding from this study and others prove that pre-hypertensive states are often unnoticed. Hence, periodic blood pressure checks, regular community screenings, and surveys for early detection and management should be encouraged to halt the progression to hypertension.

Most national and global studies, reported low awareness of hypertension status^{13,14,22-24,30,39}. However, the majority of the hypertensives in this study (73%) were aware of their status. In the studies

conducted in China and Seychelles Island, awareness rate was 45% and 50% respectively^{35,40}. Also, the previously diagnosed hypertensives checked their blood pressures more regularly than the other groups. The awareness of their hypertensive states and knowledge of possible complications may be the reason for the regular blood pressure check.

A good proportion of the hypertensives were on antihypertensive medications, however, adherence was poor and many had uncontrolled blood pressure. This finding was nearly similar to what was obtained in Zimbabwe and some studies in Nigeria^{29,31,39,40}. Erhun et al. and Daniel et.al., in their studies, reported few hypertensives on treatment with good blood pressure control^{24,36}. As documented in the American heart association guideline, about one-third of the hypertensives are on antihypertensive and over half had uncontrolled blood pressure. Many factors including underestimation of the impact of drugs, the cost of treatment have been documented as some of the reasons for poor compliance and control. Older persons are at greater risk of developing hypertension as seen in this study and reported in other studies. However, the condition has increasingly impacted all age groups. Dietary pattern changes — moving from home meals to fast foods, obesity, and stress of urbanization seen among all ages may be the major contributing factors. Males showed a higher risk of developing hypertension than females; similar to the reports in most studies^{13,14,32,36-40}.

However, Adeniran *et. al*, reported more hypertensives in females than males²⁸. This sex differential may be due to hormonal differences and other factors like greater stress burden on men as breadwinners and better access to healthcare services by females.

Although most participants have heard of hypertension, only a few had reasonably good knowledge of the disease. A large proportion had poor knowledge, especially regarding its risk factors and complications. The low knowledge as reported in this study is comparable to other studies across

Nigeria and other countries^{8,24}. In Thailand and Zimbabwe studies, knowledge was poor. In the Seychelles Island and Sri-lanka studies, knowledge was good but attitude and practices were poor. Poor knowledge continues to remain a major hindrance in curbing the menace of hypertension, especially as hypertension is usually asymptomatic.

Previous observational studies have shown the protective effect of exercise and physical activity against hypertension and its complications^{41,42}. This study also demonstrated the benefit with a good proportion of those that engaged in regular physical exercise and active lifestyle having a 53.3% risk reduction of developing high blood pressure. While the mechanism for the blood pressure lowering effect of physical activity is not very clear, it is thought to be related to modulation of the autonomic nervous system, renin-angiotensin-aldosterone system, cardiac output, peripheral vascular resistance, insulin resistance, and endothelial function⁴³.

There was no statistically significant difference in the blood pressure of the cohort with reference to alcohol consumption, likely because there were no heavy drinkers' in the study population. Recent studies suggested that moderate consumption of alcohol (<30g of ethanol) is associated with a reduction in blood pressure and consequently cardiovascular events^{44,45}. While heavy drinking (>30g of ethanol) is associated with elevation of cardiovascular risk as well as blood pressure, affecting systolic more than diastolic blood pressure^{46,47}. Nevertheless, it is always an uphill task to get unequivocal outcomes given the difficulty associated with getting accurate data on the type, pattern, and quantity of alcohol consumption with a lot of confounding socioeconomic and other lifestyle behaviour can't be adequately reflected in statistical models.

Cigarette smokers showed a higher (2.21 times) risk of developing higher blood pressure compared to non-smokers in this study. This finding was similar to a study carried out in Ukraine, which reported more hypertensive among smokers compared to

never-smokers and former smokers of less than twenty years' duration⁴⁸. Paradoxically some studies actually found lower blood pressure values among habitual smokers when compared with never-smokers and former-smokers^{49,50}. Studies carried out among female smokers in Europe and the Health Survey of England database also came to a similar conclusion^{50,51}. The effects of cigarette smoking on blood pressure are complicated. It is extensively documented that tobacco smoking acutely increases blood pressure and the risk of severe hypertension and its complications. It also accelerates atherothrombotic processes on chronic usage through possibly deleterious effects on endothelial function, inflammation, and thrombosis. This study also showed a positive correlation between BMI, waist circumference, and blood pressure. Hypertensives had a higher mean BMI and waist circumference than normotensives. This is similar to many studies that have also documented these relationships^{52,53}.

A popular quotation by scientists; is "genes are like loaded guns but lifestyle pulls the trigger"⁵⁴. Genetic makeups alone are not enough for disease development. Simple changes in lifestyle and health promotion can prevent hypertension. Although; genetic predisposition to hypertension is non-modifiable, environmental factors are modifiable and herein lay the potential to prevent hypertension.

CONCLUSION/RECOMMENDATION

This study area has progressive increase in the prevalence of systemic hypertension. A substantial number of persons found that they are hypertensive for the first time. There is a high level of awareness of the disease without a concomitant deeper understanding of the associated risk factors and complications. Respondents practice varying treatment options including the use of orthodox antihypertensive medications and herbal preparation as well as a combination of both. Nevertheless, compliance with orthodox medication was poor, with only a few having optimal BP control.

Therefore measures need to be put in place to help increase the knowledge and awareness of systemic hypertension at the community level where most of

the respondents' domicile, to stem the rising tide of systemic hypertension and its attendant complications. It is important that people undergo frequent blood pressure appraisal, especially those with relevant risk factors.

REFERENCE

1. World Health Organization. World Health Statistics 2020: Monitoring Health for SDGs. <https://www.who.int/data/gho/whs-2020-visual-summary>. Accessed date 13 May 2021 at 0051hours.
2. Dorans KS, Mills KT, Liu Y, He J. Trends in prevalence and control of hypertension according to the 2017 American college of cardiology/American heart association (ACC/AHA) guideline. *J Am Heart Assoc* 2018; 7(11): e008888. doi:10.1161/JAHA.118/008888.
3. Arima H, Barzi F, Chalmers J. Mortality patterns in hypertension. *J Hypertens* 2011;29Suppl 1:S3-7.
4. World Health Organization. A global brief on hypertension world health day, silent killer, global public health crisis. <http://www.who.int/publications/items/a-global-brief-on-hypertension-silents-killer-global-public-health-crisis-world-health-day-2013>. Accessed date 28 July 2021 1155hours.
5. Kifle ZD, Adugna M, Chanig, Mohammed A. Prevalence and associated factors for hypertension complications among hypertensive patients at University of Gondar Comprehensive Specialized Referral Hospitals. *Clinical Epidemiology and Global Health*.2022;13(100951):ISSN 2213-3984.
6. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the global burden of disease study 2010. *Lancet* 2012;380(9859):2224-60.
7. Mills KT, Stefanescu A, He J. The global epidemiology of hypertension. *Nat Rev Nephrol* 2020;16(4):223-37.

8. Ofili MI, Ncama BP, Sartorius B. Hypertension in rural communities in Delta State, Nigeria: prevalence, risk factors and barriers to health care. *Afr J Prim Health Care Fam Med* 2015;49(1):875-81.
9. Vijver SV, Akinyi H, Oti S, Olajide A, Agyemang C, Aboderin I, et al. Status report on hypertension in Africa-Consultative review for the 6th session of African union conference of ministers of health on NCDS. *Pan Afr Med J* 2013;16:38
10. Guwatudde D, Nankya-Mutyoba J, Kalyesubula R, Laurence C, Adebamowo C, Ajayi IO, et al. The burden of hypertension in sub-Saharan Africa: a four-country cross sectional study. *BMC Public Health* 2015;15(1):1211.
11. Dzau VJ, Balatbat CA. Future of hypertension: the need for transformation. *Hypertension* 2019;74(3):450-57.
12. Ogah OS, Okpechi I, Chukwuonye II, Akinyemi JO, Onwubere BJ, alase AO, et al. Blood pressure, prevalence of hypertension related complications in Nigerian Africans: a review. *World J Cardiol* 2012;4(12):327-40.
13. Okubadejo NU, Ozoh OB, Ojo OO, Akinkugbe AO, Odeniyi IA, Adegoke O, et al. Prevalence of hypertension and blood pressure profile amongst urban-dwelling adults in Nigeria: a comparative analysis based on recent guideline recommendations. *Clin Hypertens* 2019;25:7.
14. Adeloye D, Basquill C, Aderemi AV, Thomson JY, Obi FA. An estimate of the prevalence of hypertension in Nigeria: a systematic review and metaanalysis. *J Hypertens* 2015;33(2):230-42.
15. World Health Organization. Hypertension key facts . 25 August 2021 . <https://www.who.int/news-room/factsheets/detail/hypertension>. Accessed date: Accessed date 28 July 2021 1150hours.
16. Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Himmelfarb CD, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American college of cardiology/American heart association task force on clinical practice guidelines. *Hypertension*. 2018;71(6):1269-324.
17. Lackland DT. Racial differences in hypertension: implications for high blood pressure management. *Am J Med Sci* 2014;348(2):135-8.
18. Benowitz NL. Antihypertensive agents. In: Katzung BG, editor. *Basic and clinical pharmacology*. 10th ed. New York: McGraw-Hill, 2007. pp159.
19. Valsan RS, Baiser A, Seshadri S, Larson MG, Kannel WB, D'Agostino RB, et al. Residual life time risk for developing hypertension in middle-aged women and men: the Framingham heart study. *JAMA* 2002;287(8):1003-10.
20. Smith WC, Lee AJ, Crombie IK, Tunstall-Pendoe H. Control of blood pressure in Scotland: the rule of halves. *BMJ*. 1990;300(6730):981-83.
21. Ulasi II, Ijoma CK, Onwubere BJ, Arodiwe E, Onodugo O, Okafor C, et al. High prevalence and low awareness of hypertension in a market population in Enugu, Nigeria. *Int J of Hypertens* 2011;2011:869675.
22. Raji YR, Abiona T, Gureje O. Awareness of hypertension and its impact on blood pressure control among elderly Nigerians: report from the Ibadan study of aging. *Pan Afr Med J*. 2017;27:190.
23. Iyalome GBS, Iyalome SI. Hypertension-related knowledge, attitudes and life-style practices among hypertensive patients in a sub-urban Nigerian community. *Journal of Public Health Epidemiol* 2010;2(4):71-7.
24. Daniel OJ, Adejumo OA, Adejumo EN, Owolabi RS, Braimoh RW. Prevalence of hypertension among urban slum dwellers in Lagos, Nigeria. *J Urban Health* 2013; 90(6):1016-25.
25. Ibekwe RU. Modified risk factors of hypertension and sociodemographic profile in Oghara, Delta State; prevalence and correlates. *Ann Med Health Sci Res* 2015;5:71-9.
26. Kabwama SN, Bahendeka SH, Wesonga R, Mutungi G, Guwatudde D. Low consumption of fruits and vegetables among adults in Uganda:

- findings from a countrywide cross-sectional survey. *Arch Public Health* 2019;77:4.
27. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, et al. National heart, lung, and blood institute joint national committee on prevention, detection, evaluation, and treatment of high blood pressure, national high blood pressure education program coordinating committee. The seventh report of the joint national committee on, prevention, detection, evaluation, and treatment of high blood pressure: the JNC report. *JAMA* 2003;289:2560-570.
 28. Adediran OS, Okpara IC, Adeniyi OS, Jimoh AK. Hypertension prevalence in an urban and rural area of Nigeria. *J Med. Med. Sci* 2013;4(4):149-54.
 29. Ekwunife OI, Udeogaranya PO, Nwatu IL. Prevalence, awareness, treatment and control of hypertension in a Nigeria population. *2010;2(7):731-35.*
 30. Akpan EE, Ekrikpo UE, Udo AIA, Bassey BE. Prevalence of hypertension in Akwa Ibom State, South-South Nigeria: rural versus urban communities' study. *International Journal of Hypertension* 2015;2015(975819):5 pages.
 31. Chimberengwa PT, Naidoo M, Cooperative Inquiry Group. Knowledge, attitudes and practices related to hypertension among residents of a disadvantage rural community in southern Zimbabwe. *PLoS ONE* 2019;14(6):e0215500.
 32. Aung MN, Lorga T, Srikrajang J, Promtingkran N, Kreuangchai S, Tonpanya W, et. al. Assessing awareness and knowledge of hypertension in an at-risk population in the Karen ethnic rural community, Thasongyang, Thailand. *Int J Gen Med* 2012(5):553-61.
 33. Muntner P, Carey RM, Gidding S, Jones DW, Taler SJ, Wright JT, et. al. Potential US population impact of the 2017 ACC/AHA high blood pressure guideline. *Circulation* 2018;137:109-18.
 34. Ajayi IO, Sowemimo IO, Akpa OM, Ossai NE. Prevalence of hypertension and associated factors among residents of Ibadan-North local government area of Nigeria. *Nig J Cardiol* 2016;13:67-5.
 35. Gao Y, Chen G, Tian H, Lin L, Lu J, Weng J, et al. Prevalence of hypertension in China: a cross-sectional study. *PLoS One* 2013;8(6):e65938.
 36. Erhun WO, Olayiwola G, Agbani EO, Omotoso NS. Prevalence of hypertension in a university community in south west Nigeria. *African Journal of Biomedical Research* 2005;8:15-19.
 37. Akinula JT, Meakin R, Umar AM, Freemantle N. Current prevalence pattern of hypertension in Nigeria: a systematic review. *PLoS One* 2015;10(10):e0140021.
 38. Osunkwo D, Mohammed A, Kamateeka M, Nguku P, Umeokonkwo CD, Abolade OS et al. Population-based prevalence and associated risk factors of hypertension among adults in Benue State, Nigeria. *Niger J Clin Pract* 2020;23(7):944-49.
 39. Odili AN, Chori BS, Nwakile IC, Okoye IC, Abdullah U, Nwegbu MN, et. al. Prevalence, awareness, treatment and control of hypertension in Nigeria: data from a nationwide survey 2017. *Global Heart* 2020;15(1):47.
 40. Aubert L, Bovet P, Gervasoni JP, Rwebogora A, Waeber B, Paccaud F. Knowledge, attitude, and practices on hypertension in a country in epidemiological transition. *Hypertension* 1998;31(5):1136-45.
 41. Cornelissen VA, Smart NA. Exercise training for blood pressure: a systematic review and meta-analysis. *J Am Heart Assoc.* 2013;2 doi:10.1161/JAHA.112.004473.
 42. Cornelissen VA, Fagard RH. Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *Hypertension* 2005;46(4):667-75.
 43. Arakawa K. Antihypertensive mechanism of exercise. *J Hypertens* 1993;11(3):223-9.
 44. Worm N, Belz GG, Stein-Hammer C. Moderate wine consumption and prevention of coronary heart disease. *Dtsch Med Wochenschr.* 2013;138(51-52):2653-57.
 45. Bos S, Grobbee DE, Boer JM, Verschuren WM, Beulens JW. Alcohol consumption and risk of

- cardiovascular disease among hypertensive women. *Eur J Cardiovasc Prev Rehabil* 2010;17:119–26.
46. Skliros EA, Papadodima SA, Sotiropoulos A, Xipnitos C, Kollias A, Spiliopoulou CA. Relationship between alcohol consumption and control of hypertension among elderly Greeks. The Nemea primary care study. *Hellenic J Cardiol* 2012;53(1):26–32.
 47. Kaplan NM. Alcohol and hypertension. *Lancet* 1995;345(8965):1588–89.
 48. Leskiv I, Povroznyk N, Mulka L, Cherevychnyk N, Dzyaman M. Effect of smoking on blood pressure control in hypertensive men. *J of Hypertens* [2015;33\(suppl 1\):e272.](#)
 49. Seltzer CC. Effect of smoking on blood pressure. *Am Heart J* 1974;87(5):558-64
 50. Masala G, Bendinelli B, Versari D, Saieva C, Ceroti M, Santagiuliana F, et al. Anthropometric and dietary determinants of blood pressure in over 7000 Mediterranean women: the European prospective investigation into cancer and nutrition-Florence cohort. *J Hypertens* 2008;26(11):2112-20.
 51. Primatesta P, Falaschetti E, Gupta S, Marmot MG, Poulter NR. Association between smoking and blood pressure: evidence from the health survey for England. *Hypertension* 2001;37(2):187-93
 52. Brown CD, Higgins M, Donato KA, Rohde FC, Garrison R, Obarzanek E, et al. Body mass index and the prevalence of hypertension and dyslipidemia. *Obes Res.* 2000;8(9):605–19.